

# Study The Fluorescence Properties For (C<sub>14</sub>H<sub>14</sub>CLN<sub>3</sub>) Molecular And Effect Of Solvent Polarity

Asmaa Satar Jeeiad Mheesin Al-Ragehey

Faculty of Physics, College of Sciences, Mustansiriyah University, Baghdad, Iraq

E-mail address : ahmed\_naji\_abd@yahoo.com

**Abstract**—The absorption and fluorescence spectra have been studied and measured for (C<sub>14</sub>H<sub>14</sub>CLN<sub>3</sub>) AF solution which dissolved in [N-N-dimethyl formamide (Dmf) , dimethyl sulphoxide (DmsO) and water] solvents as a function of concentration and solvent at excitation wavelength ( $\lambda_{ex}$ = 414nm) . Have been calculated peaks the fluorescence spectra of (C<sub>14</sub>H<sub>14</sub>CLN<sub>3</sub>) solution dissolved in N-N-dimethyl formamide (Dmf) and dimethyl sulphoxide (DmsO) at concentration ( $1 \times 10^{-5}$ ) M, and found that it is equal to ( $\lambda_{max}$ = 497 ) , ( $\lambda_{max}$ = 501) and ( $\lambda_{max}$ = 505), respectively. Also been calculated the value stokes shift of these peaks found to be equal to (1292.5) , (1316.5) and (2469.7) , respectively. Where we note that (stokes shift) shifted peak of fluorescence spectra towards the longer wavelength (redshift) increases with increase solvent polarity. The increase of solvents polarity lead to decrease in the intensity of the fluorescence spectra. All measurement were at room temperature .

**Keywords**—Absorption and fluorescence spectra , Acriflavine molecular, Concentration , Stokes shift and Redshift.

## 1.Introduction

Many studies have been reported for the (C<sub>14</sub>H<sub>14</sub>CLN<sub>3</sub>) molecule, such as M. Maeda in 1984 has been used the acriflavine as active medium for dye lasers [1]. Carlos M. Previtali in 1995 investigated the solvent effects on the rate constant and activation parameters of several intermolecular electron transfer reactions [2]. Vijay K. Sharma *et.al.* in 2008 study that has been shown the magnitude of the Stokes shift (frequency shifts in absorption and fluorescence spectra) is observed on changing the solvents and further has been used to calculate experimentally the dipole moments (ground state and excited state) of acriflavine and acridine orange dye molecules [3]. M Lee lavathi *et.al.* in 2011 investigated the acriflavine lotion as topical antiseptic solution yellow or orange in colour, mainly used for minor wounds, burns, and infected skin. Although used in dilution 0.1% for medical purposes, this agent has been documented to produce potential skin itchiness, irritation or burning sensation upon contact. Nevertheless, it is still widely used for wound dressing by health professionals and remains a popular topical antiseptic agent purchased

over- the counter [4]. It is well known that the loss of energy of an excited molecule in a solution, both by radiative and non-radiative processes characteristically depend upon the molecular structure and the environment of the molecules. Therefore, not only the structure of solute molecules but the solvent also plays an important role in determining its radiative and non- radiative properties

A change for the type of a solvent when it produces alteration in both the excitation and fluorescence wavelength indicates that the solvent, in some way, is able to interact with the solute in the ground state. However, when solvent produces a shift in fluorescence wavelength only, it indicates an interaction between solvent and solute molecule, in excited state. Therefore , when a molecule is excited, its dipole moment gets changed and it remains no more in equilibrium with its immediate environment. As a molecule relaxes and is in equilibrium with its surroundings, some energy is dissipated in the form of heat energy and the fluorescence emission wavelength gets shifted according to Frank - Condon principle [5]. There are many processes bimolecular which commonly compete with fluorescence emission and internal quenching in solutions, and thereby modify the fluorescence characteristics. This processes are collision impurity quenching, concentration quenching ,energy transfer quenching and self-absorption quenching [6].

The self-absorption quenching is in principle, an increase in the concentration of the fluorescence solute in a given material should be accompanied by an increase in the emitted light intensity. This is due to the corresponding increase in the absorption efficiency. However, such behavior only occurs up to a certain critical concentration of the fluorescence solute. Above this concentration, the fluorescence intensity starts to decrease. This process is known as concentration quenching of fluorescence [7]. The interactions responsible for general solvent effects are best understood by derivation of the Lippert equation. This equation can be written as follows:

$$hc \Delta \bar{\nu} = hc(\bar{\nu}_a - \bar{\nu}_f) = \frac{2\Delta f}{a^3} (\mu_E - \mu_G)^2 + constant. (1)$$

Where:  $\Delta \bar{\nu}$  is the frequency shift in  $\text{cm}^{-1}$  between absorption and emission,  $a$  is the cavity radius, and  $\mu_E$  and  $\mu_G$  are the excited and ground state dipole

moments, respectively. The  $\Delta f$  is called orientation polarizability, one can find this parameter as in the following equation 2 [8].

$$\Delta f = \frac{\epsilon - 1}{2\epsilon + 1} - \frac{n^2 - 1}{2n^2 + 1} \quad (2)$$

Where :  $\epsilon$  is the dielectric constant,  $n$  is the refractive index.

## 2.Experimental:

### 2-1 materials:

The Acriflavine (AF) is derived from (acridine family) as be shown in figure 1. It dissolved in different polarity solvent like [*N-N*-dimethyl formamide (Dmf), dimethyl sulphoxide (DmsO) and water] as solutions, prepared at concentration [ $1 \times 10^{-5}$ ] M. (AF) purchased from Uma Company imported from India.




Fig.(1): Chemical formula of Acriflavine ( $C_{14}H_{14}CLN_3$ ) [3].

### 2-2 Absorption and Fluorescence measurements:

Absorption spectrum was measured by a UV-V is spectrophotometer CARY 100 Conc and made by OPTIMA INC January 2009.


Fluorescence spectrum was record by using spectrofluorophotometer kind of RF-5301 pc Shimadzu sample were mounted cubic cell of quartz dimensions ( $1 \times 1 \times 5$ ) cm at angle 90 with incident beam. This optical geometry was chosen to eliminate the effect of scattered incident radiation and the self-absorption phenomena. The instrument computerized and operates in the wavelength range 220-900 nm. The fluorometer has dedicated computer which control instrumental operating excitation and emission wavelength, scan, monochromator slit width, detector parameter and the acquisition of spectral data. The exiting wavelength  $\lambda_{ex}$  (414 nm) is used in the measurements and all measurements were at room temperature performances.

## 3.Results and Discussion:


In this paper, the fluorescence spectra of ( $C_{14}H_{14}CLN_3$ ) solution dissolved in different polarity solvent like [*N-N*-dimethyl formamide (Dmf), dimethyl sulphoxide (DmsO) and water], are studied with concentration [ $1 \times 10^{-5}$ ] M and the effect of polarity on the fluorescence spectrum.

Figure 2 and figure 3 shows the absorption spectra and the fluorescence spectra of acriflavine


solutions in different solvent polarity, respectively . From the figure 3 the fluorecence spectra appears as structure-less with peak located at wavelength in (*N-N*-dimethyl formamide (Dmf)  $\lambda_{max}$ = 497 nm), (dimethyl sulphoxide (DmsO)  $\lambda_{max}$ = 501 nm) and (water  $\lambda_{max}$ = 505 nm).



C) AF-Water






a) AF-DMF



b) AF-DMSO


Fig.(2): Absorption spectra of Acriflavine in [a] *N-N*-dimethyl formamide (Dmf) , b)dimethyl sulphoxide (DmsO) and c)water] at concentration.



**Fig.(3): Fluorescence spectra of Acriflavine in [a) N-N-dimethyl formamide (Dmf) , b)dimethyl sulphoxide (DmsO) and c)water] at concentration  $1 \times 10^{-5}$  M.**

From the figure 4 can observe the effected of the solvent polarity on the fluorescence spectra, where the fluorescence intensity decreases, when polarity of solvent are increased because the molecule when absorbed the light and transitions to the excited electronic state, the molecule electronic configuration is changed and hence solute-solvent molecule interacts dipole-dipole [10]. From the figure 4, the acriflavine dye , It suggests that it is a locally excited intra molecular charge transfer ICT state. Amino group in the mero-phenyl ring is protonated to give

high fluorescent and as a result, twisted ICT (TICT) states are not formed in case of acriflavine dye or interacting between solute with molecule solvent as dipole-dipole interacts .[11]




**Fig.(4): fluorescence spectra of acriflavine solution in [ N-N-dimethyl formamide (Dmf) , dimethyl sulphoxide (DmsO) and water] at concentration  $1 \times 10^{-5}$  M.**

From the last figure, one can observe that, when the polarity was increased the fluorescence spectrum shifted to a long wavelength red shift and decrease in the intensity of fluorescence. This shift value is dependent on the dielectric constant  $\epsilon$  value of solvent and on the difference between dipole moment values of solvent [12], as shown in Table 1.

**Table (1): photo-physical parameters of acriflavine in different solvents at room temperature.**

solvent	$\lambda_a$ (nm)	$\lambda_f$ (nm)	$\nu_s^-$ ( $\text{cm}^{-1}$ )	$\nu_f^-$ ( $\text{cm}^{-1}$ )	$\nu_s^- - \nu_f^-$ ( $\text{cm}^{-1}$ )	$\frac{\epsilon_0 + \epsilon_f}{2}$ ( $\text{cm}^{-1}$ )	$\Delta f$
Dmf	467	497	21413.2	20120.7	1292.5	20766.9	0.275
DmsO	470	501	21276.5	19960	1316.5	20618.25	0.263
Water	449	505	22271.7	19801.9	2469.7	21036.8	0.320



**Fig.(5): the relation between the stokes shift value and orientation polarizability  $\Delta f$  acriflavine molecule dissolved in [ N-N-dimethyl formamide (Dmf) , dimethyl sulphoxide (Dmso) and water]**

**at concentration [ $1 \times 10^{-5}$  M].**

#### 4. Conclusions

Throughout the study of acriflavine AF dye dissolved in [N-N-dimethyl formamide (Dmf) , dimethyl sulphoxide (Dmso) and water] with a concentration [ $1 \times 10^{-5}$ ] M the following conclusions are reached to :

1- The fluorescence intensity decreases with the increase of solvent polarity because of the formation of exciplexes complex or intra molecular charge-transfer ICT phenomenon.

2- The red shift of fluorescence was increased with the increasing of solvent polarity.

#### References

1- M. Maeda, Laser Dyes, Academic Press, New York,(1984).

2- C. M. Previtali, "Solvent effects on intermolecular electron transfer processes", J. Appl. Chem, Vol. 67 (1995) 127-134.

3- Vijay K. Sharma , P.D. Sahare , Ramesh C. Rastogi , S.K. Ghoshal and D. Mohan; "Excited state characteristics of acridine dyes: acriflavine and acridine orange", J. Elsevier Science B.V. ,Vol. 59 (2008) 1799–1804 .

4- M Lee Lavathi , YY Le , H Tohid and AH Hasliza , " Contact dermatitis presenting as non-healing wound: case report" , J. Asia Pacific Family Medicine,Vol. 10 (2011) 1-3.

5- D.M. Hercules (Ed.), "Fluorescence and Phosphorescence Analysis Principals and Applications", Interscience, (2013).

6- J.N. Demas and G.A. Crosby, "The measurement of photoluminescence quantum yields", J. Phys. Chem., Vol.75 (2010) 991-1031.

7- B. Valeur, "Molecular fluorescence: principles and applications", Wiley-VcH Verlag GmbH, (2011).

8- A. S. R. Koti and N. Periasamy, "Solvent Exchange in Excited-State Relaxation in Mixed Solvents", Journal of Fluorescence, Vol.10 (2014) 177-184.

9- P. L. Muir o and P. R. Callis," Solvent Effects on the Fluorescence Quenching of Tryptophan by Amides via Electron Transfer. Experimental and Computational Studies", J. Phys. Chem., (2008).

10- B.M. Krasontskin, B.M. Bolotin, Organic Luminescent Materials, Weinheim VCH, (2012).

11- J. Petit, M. Denis-Gay and Marie-H el ene Ratinaud, " Assessment of fluorochromes for cellular structure and function studies by flow cytometry ", Biol Cell, Vol. 78 (2013) 1-13 .