

Estimates for a Hyperbolic Equation with no Dissipation on R^N .

Perikles Papadopoulos and Niki-Lina Matiadou

Department of Electronics Engineering,
 Piraeus University of Applied Sciences, Greece
ppapadop@teipir.gr, lmatiadou@yahoo.gr

Abstract— We discuss the energy decay estimates and the local existence results of the solutions for the nonlocal hyperbolic problem with no dissipation

$u_{tt} + \phi(x) \|\nabla u(t)\|^2 (-\Delta)u = 0$, $x \in R^N$, $t \geq 0$, with initial conditions $u(x, 0) = u_0(x)$ and $u_t(x, 0) = u_1(x)$, in the case where $N \geq 3$, and $(\phi(x))^{-1} = g(x)$ is a positive function lying in $L^{N/2}(R^N) \cap L^\infty(R^N)$. We prove that we have unique local solution for our problem in the case when the initial energy is small.

Keywords— Quasilinear Hyperbolic Equations, Unbounded Domains, Generalized Sobolev Spaces.

I. Introduction

In this work we study the following degenerate wave equation with no dissipation

$$(1.1) \quad u_{tt} + \phi(x) \|\nabla u(t)\|^2 (-\Delta)u = 0,$$

$$(1.2) \quad u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x),$$

$x \in R^N, t \geq 0$, with initial conditions u_0, u_1 in appropriate function spaces, $N \geq 3$. The case of $N=1$, equation (1.1) describes the nonlinear vibrations of an elastic string. Throughout the paper we assume that the function ϕ and $g: R^N \rightarrow R$ satisfy the following condition

$$(G) \quad \phi(x) > 0, \quad \text{for all } x \in R^N \quad \text{and} \\ (\phi(x))^{-1} = g(x) \in L^{N/2}(R^N) \cap L^\infty(R^N).$$

This class will include functions of the form $\phi(x) \square c_0 + \varepsilon |x|^a, \varepsilon > 0$ and $a > 0$, resembling phenomena of slowly varying wave speed around the constant speed c_0 . Many results treat the case of $\phi(x) = \text{constant}$ (in bounded or unbounded domains). It must be noted, that this case is proved to be totally different from the case of $\phi(x) \rightarrow c_\pm > 0$, as $x \rightarrow \pm\infty$ (see [8]).

The original equation is

(1.3)

$$ph \frac{\partial^2 u}{\partial t^2} + \delta \frac{\partial u}{\partial t} = \left\{ p_0 + \frac{Eh}{2L} \int_0^L \left(\frac{\partial u}{\partial x} \right)^2 \right\} \frac{\partial^2 u}{\partial x^2} + f \quad \text{for } 0$$

$< x < L, t \geq 0$, where $u = u(x, t)$ is the lateral displacement at the space coordinate x and the time t , E the Young modulus, p the mass density, h the cross-section area, L the length, p_0 the initial axial tension, δ the resistance modulus and f the external force. When $p_0 = 0$ the equation is considered to be of *degenerate type* and the equation models an unstretched string or its higher dimensional generalization. Otherwise it is of *nondegenerate type* and the equation models a stretched string or its higher dimensional generalization. When $\delta = f = 0$, the equation was introduced by G. Kirchhoff [12] in the study of oscillations of stretched strings and plates. That's why equation (1.3) is called the *Kirchhoff string*.

In the case treated here the problem becomes complicated because the equation does not give rise to compact operators. The homogeneous Sobolev spaces combined with equivalent weighted L^p spaces, is the appropriate space to overcome these difficulties. In our paper we assume that $f(u) = 0$ (we have no external force), in order to study the behavior of the solutions for this kind of equations. This case is rather interesting in the class of the homogeneous Sobolev spaces as we will study. In the case of *bounded domain*, when $\delta = 0$ and $f \neq 0$, the global existence is rather well studied in the class of analytic function spaces (e.g. see [5]). H. Crippa [3] has proved local in time solvability in the class of usual Sobolev spaces. A. Arosio and S. Garavaldi [1] have shown the existence of a unique local solution in the case of mildly degenerate type. For $\delta \geq 0$ and $f(u) = 0$, in the degenerate case, the global existence of solutions has been shown by K. Nishihara and Y. Yamada [16], when the initial data are small enough. When $\delta > 0$ and $f(u) = 0$, M. Nakao [14] has derived decay estimates for the solutions. In particular, T. Kobayashi [13] constructed a unique weak solution using a Faedo-Galerkin method for a quasilinear wave equation with strong dissipation (see also [4, 15]). K. Nishihara [17] has derived a decay estimate from

below of the potential of solutions. In the case of $\delta \geq 0$ and $f \neq 0$, M. Hosoya and Y. Yamada [7] have studied the non-degenerate case with linear dissipation and proved the global existence of a unique solution under small initial data. R. Ikehata [9] has shown that for sufficiently small initial data, global existence can be obtained, even when the influence of the source terms is stronger than that of the damping terms.

In the case of *unbounded domains*, P. 'Ancona and S. Spagnolo [6] have shown the global existence of a unique C^∞ solution for the non-degenerate type with small C_0^∞ data. G.Todorova [21] studied the global existence and nonexistence of solutions both in the bounded and unbounded domain cases with nonlinear damping and small enough C_0^∞ initial data.

Finally, N. Karahalios and N. Stavrakakis [10]-[11] have proved global existence and blow-up results for some semilinear wave equations with weak damping on all R^N .

The presentation of this paper has as follows: In Section 2 we discuss properties of the homogeneous Sobolev space $D^{1,2}(R^N)$ and some weighted L^p spaces, in order to overcome difficulties of non-compactness arising from the unboundedness of the domain. In Section 3, we show the existence of a unique local weak solution and we obtain energy decay estimates for the problem (1.1)-(1.2) with

$$(u_0, u_1) \in D^{1,2}(R^N) \times L_g^2(R^N),$$

when the initial energy $E(u_0, u_1)$ which corresponds to the problem, is non-negative and small.

Notation: We denote by B_R the open ball of R^N with center 0 and radius R. Sometimes for simplicity we use the symbols C_0^∞ , $D^{1,2}$, L^p , $1 \leq p \leq \infty$, for the spaces $C_0^\infty(R^N)$, $D^{1,2}(R^N)$, $L^p(R^N)$, respectively; $\|\cdot\|_p$ for the norm $\|\cdot\|_{L^p(R^N)}$, where in case of $p = 2$ we may omit the index.

II. Preliminary Results

In this section, we briefly mention some facts, notation and results, which will be used later in this paper. The space $D^{1,2}(R^N)$ is defined as the closure of $C_0^\infty(R^N)$ functions with respect to the energy norm $\|u\|_{D^{1,2}} = \left(\int_{R^N} |\nabla u|^2 dx \right)^{1/2}$. It is known that

$$D^{1,2}(R^N) = \left\{ u \in L^{\frac{2N}{N-2}}(R^N) : \nabla u \in (L^2(R^N))^N \right\}$$

and $D^{1,2}(R^N)$ is embedded continuously in $L^{\frac{2N}{N-2}}(R^N)$, that is, there exists $k > 0$ such that

$$(2.1) \quad \|u\|_{L^{\frac{2N}{N-2}}(R^N)} \leq k \|u\|_{D^{1,2}}.$$

We shall frequently use the following generalized version of Poincaré's inequality

$$(2.2) \quad \int_{R^N} |\nabla u|^2 dx \geq \alpha \int_{R^N} gu^2 dx,$$

for all $u \in C_0^\infty$ and $g \in L^{\frac{N}{2}}$, where

$$\alpha = k^{-2} \|g\|_{L^{N/2}}^{-1} \quad (\text{see [2, Lemma 2.1]}).$$

It is shown that $D^{1,2}(R^N)$ is a separable Hilbert space. The space $L_g^2(R^N)$ is defined to be the closure of

$C_0^\infty(R^N)$ functions with respect to the inner product

$$(2.3) \quad (u, v)_{L_g^2(R^N)} = \int_{R^N} guv dx.$$

It is clear that $L_g^2(R^N)$ is a separable Hilbert space. The following Lemmas will be proved to be useful in the sequel. For the proofs we refer to [11], (we note that g is a positive function).

Lemma 2.1 Let $g \in L^{N/2}(R^N) \cap L^\infty(R^N)$. Then the embedding $D^{1,2} \subset L_g^2$ is compact.

Lemma 2.2 Let $g \in L^{\frac{2N}{2N-pN+2p}}(R^N)$. Then the following continuous embedding $D^{1,2} \subset L_g^p$ is valid, for all $1 \leq p \leq 2N/(N-2)$.

Remark 2.3 The assumption of Lemma 2.2 is satisfied under the hypothesis (G), if $p \geq 2$.

Lemma 2.4 Let g satisfy condition (G). If $1 \leq q < p < p^* = 2N/(N-2)$, then the following weighted inequality

$$(2.4) \quad \|u\|_{L_g^p} \leq C_0 \|u\|_{L_g^q}^{1-\theta} \|u\|_{D^{1,2}}^\theta,$$

is valid, for all $\theta \in (0, 1)$, for which

$$1/p = (1-\theta)/q + \theta/p^*, \text{ and } C_0 = k^\theta.$$

To study the properties of the operator $-\phi\Delta$, we consider the equation

$$(2.5) \quad -\phi(x)\Delta u(x) = \eta(x), x \in R^N,$$

without boundary conditions. Since for every $u, v \in C_0^\infty$ we have

$$(2.6) \quad (-\phi\Delta u, v)_{L_g^2} = \int_{R^N} \nabla u \nabla v dx,$$

we may consider equation (2.5) as an operator equation of the form

$$(2.7) \quad A_0 u = \eta, A_0 : D(A_0) \subseteq L_g^2 \rightarrow L_g^2, \eta \in L_g^2.$$

Relation (2.6) implies that the operator $A_0 = -\phi\Delta$ with domain of definition $D(A_0) = C_0^\infty$, is symmetric. From (2.2) and equation (2.6) we have that

$$(2.8) \quad (A_0 u, u)_{L_g^2} \geq \alpha \|u\|_{L_g^2}^2, \text{ for all } u \in D(A_0).$$

So the operator $A_0 = -\phi\Delta$ is a symmetric, strongly monotone operator on L_g^2 . Hence, Friedrich's extension theorem [22, Theorem 19.C] is applicable. The energetic scalar product given by (2.6) is

$$(u, v)_E = \int \nabla u \nabla v dx$$

and the energetic space is the completion

of $D(A_0)$ with respect to $(u, v)_E$. It is obvious that the energetic space X_E is the homogeneous Sobolev space $D^{1,2}$. The energetic extension $A_E = -\phi\Delta$ of A_0 , (2.9) $-\phi\Delta : D^{1,2} \rightarrow D^{-1,2}$,

is defined to be the duality mapping of $D^{1,2}$. We define $D(A)$ to be the set of all solutions of equations (2.5), for arbitrary $\eta \in L_g^2$. Friedrich's extension A of A_0 is the restriction of the energetic extension A_E to the set $D(A)$. The operator $A = -\phi\Delta$ is self-adjoint and therefore graph-closed. Its domain $D(A)$, is a Hilbert space with respect to the graph scalar product

$$(u, v)_{D(A)} = (u, v)_{L_g^2} + (Au, Av)_{L_g^2}, \quad \text{for all } u, v \in D(A).$$

The norm induced by the scalar product is

$$\|u\|_{D(A)} = \left\{ \int_{R^N} g |u|^2 dx + \int_{R^N} \phi |\Delta u|^2 dx \right\}^{1/2},$$

which is equivalent to the norm

$$\|Au\|_{L_g^2} = \left\{ \int_{R^N} \phi |\Delta u|^2 dx \right\}^{1/2}. \text{ So we have}$$

established the evolution triple

$$(2.10) \quad D(A) \subset D^{1,2} \subset L_g^2 \subset D^{-1,2},$$

where all the embeddings are dense and compact. Finally, for later use, it is necessary to remind that the eigenvalue problem

$$(2.11) \quad -\phi(x)\Delta u = \mu u, \quad x \in R^N,$$

has a complete system of eigensolutions $\{w_n, \mu_n\}$ satisfying the following properties

$$(2.12) \quad \begin{cases} -\phi\Delta w_j = \mu_j w_j, j = 1, 2, \dots, w_j \in D^{1,2}, \\ 0 < \mu_1 \leq \mu_2 \leq \dots, \mu_j \rightarrow \infty, \text{ as } j \rightarrow \infty. \end{cases}$$

In order to clarify the kind of solutions we are going to obtain for the problem (1.1)-(1.2), we give the definition of the weak solution for this problem.

Definition 2.5 A weak solution of the problem (1.1)-(1.2) is a function u such that

$$u \in L^2[0, T; D(A)], u_t \in L^2[0, T; D^{1,2}],$$

$$(i) \quad u_{tt} \in L^2[0, T; L_g^2],$$

(ii) for all $v \in C_0^\infty([0, T] \times (R^N))$, satisfies the generalized formula

$$(2.13) \quad \int_0^T (u_{tt}(\tau), v(\tau))_{L_g^2} d\tau + \int_0^T \left(\|\nabla u(\tau)\|^{2\gamma} \int_{R^N} \nabla u(\tau) \nabla v(\tau) dx \right) d\tau = 0$$

and (iii) satisfies the initial conditions

$$u(x, 0) = u_0(x) \in D^{1,2}, u_t(x, 0) = u_1(x) \in L_g^2.$$

III. Existence Results and Energy Decay Estimates

In order to obtain a local existence result for the problem (1.1)-(1.2), we need information concerning the solvability of the corresponding non-homogeneous linearized problem restricted in the sphere $B_R : (3.1)$

$$u_{tt} - \phi \|\nabla v\|^2 \Delta u = 0, (x, t) \in B_R \times (0, T),$$

$$u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), x \in B_R,$$

$$u(x, t) = 0, (x, t) \in \partial B_R \times (0, T),$$

$$v \in C(0, T; D^{1,2}) \text{ and } v_t \in C(0, T; L_g^2).$$

Proposition 3.1 Assume that $u_0 \in D^{1,2}, u_1 \in L_g^2$ and $N \geq 3$, then the linear wave equation (3.1) has a unique solution such that

$$u \in C(0, T; D^{1,2}) \text{ and } u_t \in C(0, T; L_g^2).$$

Proof. The proof follows the lines of [11, Proposition 3.1]. The Galerkin method is used, based on the information taken from the eigenvalue problem (2.11).

Next, we will prove the following Theorem

Theorem 3.2 We assume that $N \geq 3$ and $u_0 \neq 0$.

$(u_0, u_1) \in D^{1,2} \times L_g^2$ and satisfy the non-degenerate condition $\|\nabla u_0\|^2 > 0$, then there exists

$T = T(\|u_0\|_{D^{1,2}}, \|\nabla u_1\|^2) > 0$, such that the problem (1.1)-(1.2) admits a unique local weak solution u satisfying:

$u \in C(0, T; D^{1,2}), u_t \in C(0, T; L_g^2)$. Moreover, if $\|\nabla u(t)\| > 0$ and $\|u(t)\|_{D^{1,2}} + \|u_t(t)\|_{L_g^2} < \infty$ for $t \geq 0$, then $T = \infty$.

Proof. For $T > 0$ and $R > 0$, we define the two parameter space of solutions

$$X_{T,R} =: \left\{ v \in C(0, T; D^{1,2}) : v_t \in C(0, T; L_g^2), \right. \\ \left. v(0) = u_0, v_t(0) = u_1, e(v(t)) \leq R^2, t \in [0, T] \right\},$$

where $e(u(t)) = \|u_t(t)\|_{L_g^2}^2 + \|u(t)\|_{D^{1,2}}^2$.

It is easy to see that $X_{T,R}$ can be organized as a complete metric space with the distance $d(u, v) =: \sup_{0 \leq t \leq T} e_1(u(t) - v(t))$, where

$e_1(v) =: \|v_t\|_{L_g^2}^2 + \|v\|_{D^{1,2}}^2$. We define the non-linear mapping S in the following way.

For every $v \in X_{T,R}, u = Sv$ is the unique solution of our problem. Using the fact that $\|\nabla u_0\| \equiv M_0 > 0$, we prove that there exist $T > 0, R > 0$ such that S maps $X_{T,R}$ into itself and S is a contraction mapping with respect to the metric $d(., .)$. By applying the Banach contraction mapping theorem, we obtain a unique solution u belonging to $X_{T,R}$. Therefore it follows from the continuity argument for wave equations that this solution u belongs to our space. For more details we refer to [18].

Next, we multiply equation (1.1) by $2gu_t$ and integrate over R^N to get

$$2 \int_{R^N} gu_{tt} u_t dx - 2 \int_{R^N} \|\nabla u(t)\|^2 \Delta u u_t dx = 0. \text{ So we get}$$

$$\frac{d}{dt} \left\{ \|u(t)\|_{L_g^2}^2 + \frac{1}{2} \|\nabla u(t)\|^4 \right\} = 0. \text{ We define the}$$

energy for our problem

$$(3.2) \quad E(t) = \|u(t)\|_{L_g^2}^2 + \frac{1}{2} \|\nabla u(t)\|^4.$$

So, we obtain the following relation

$$(3.3) \quad \frac{d}{dt} E(t) = 0.$$

We integrate the previous equation in $[0, t]$ to get the following

$$\int_0^t \frac{d}{dt} E(t) dt = 0, \quad E(t) - E(0) = 0, \text{ and finally}$$

$$(3.4) \quad E(t) = E(0).$$

Next, we multiply relation (3.2) by $2ug$ and integrate over R^N to get

$$2uu_{tt}g(x) - 2\phi(x)g(x)\|\nabla u(t)\|^2 = 0 \text{ and}$$

$$(3.5) \quad \int_{R^N} 2uu_{tt}g dt - \int_{R^N} 2\|\nabla u\|^2 \Delta u u dt = 0. \text{ On the}$$

other hand we have the following relation $(uu_t)' = u_t u_t + uu_{tt}$. Thus we get

$$(3.6) \quad uu_{tt} = (uu_t)' - u_t^2, \text{ and}$$

$$\int_{R^N} 2uu_{tt}g dt = \frac{d}{dt} \int_{R^N} 2guu_t dt - \int_{R^N} 2gu_t^2 dt.$$

Then we obtain

$$(3.7) \quad \int_{R^N} 2uu_{tt}g dt = \frac{d}{dt} 2(u, u_t)_{L_g^2} - 2\|u_t\|_{L_g^2}^2.$$

Using relations (3.6) and (3.7), relation (3.5) becomes

$$(3.8) \quad \frac{d}{dt} 2(u, u_t)_{L_g^2} - 2\|u_t\|_{L_g^2}^2 - \int_{R^N} 2\|\nabla u\|^2 \Delta u u dt = 0$$

where we have that

$$(3.9) \quad - \int_{R^N} 2\|\nabla u\|^2 \Delta u u dt = 2\|\nabla u\|^2 \|\nabla u\|^2 \\ = 2\|\nabla u(t)\|^4,$$

where we used the relation $-\int_{R^N} \Delta u u dt = \|\nabla u\|^2$.

Next, using relation (3.9), we obtain from relation (3.8) the following

$$\frac{d}{dt} 2(u, u_t)_{L_g^2} - 2\|u_t\|_{L_g^2}^2 + 2\|\nabla u\|^4 = 0. \text{ Thus we}$$

get the following equality

$$(3.10) \quad \frac{d}{dt} \left\{ 2(u, u_t)_{L_g^2} \right\} + 2\|\nabla u\|^4 = 2\|u_t\|_{L_g^2}^2.$$

We integrate relation (3.10) in $[0, t]$ and we get

$$2 \frac{d}{dt} \int_0^t (u(t), u_t(t))_{L_g^2} dt + 2 \int_0^t \|\nabla u(t)\|^4 dt = 2 \int_0^t \|u_t(t)\|_{L_g^2}^2 dt.$$

So, we have that

$$2(u, u_t)_{L_g^2} - 2(u_0, u_1)_{L_g^2} + 2 \int_0^t \|\nabla u(t)\|^4 dt = 2 \int_0^t \|u_t\|_{L_g^2}^2 dt.$$

Thus, we obtain the following estimate

$$(3.11) \quad 2 \int_0^t \|\nabla u\|^4 dt \leq 2(u_0, u_1)_{L_g^2} \\ + 2\|u\|_{L_g^2} \|u_t\|_{L_g^2} + 2 \int_0^t \|u_t\|_{L_g^2}^2 dt.$$

From relations (3.2) and (3.4), we get the following equality

$$\|u_t(t)\|_{L_g^2}^2 + \frac{1}{2} \|\nabla u(t)\|^4 = E(0). \text{ Thus we have}$$

that

$$(3.12) \quad \|u_t(t)\|_{L_g^2}^2 \leq E(0) \quad \text{and}$$

$$(3.13) \quad \frac{1}{2} \|\nabla u\|^4 \leq E(0) \Rightarrow \|\nabla u\|^2 \leq (2E(0))^{1/2}.$$

We obtain from relation (3.12) that

$$\begin{aligned} & 4 \int_0^t \|\nabla u\|^4 dt \leq + 2(u_0, u_1)_{L_g^2} \\ & + 2 \|u\|_{L_g^2} \|u_t\|_{L_g^2} + 2 \int_0^t \|u_t\|_{L_g^2}^2 dt + \\ & 2 \int_0^t \|\nabla u\|^4 dt \leq 2(u_0, u_1)_{L_g^2} + E(0) + E(0). \end{aligned}$$

So, we have (using Young's inequality)

$$\begin{aligned} & 4 \int_0^t \|\nabla u\|^4 dt \leq 2(u_0, u_1)_{L_g^2} + 2E(0) \\ & \leq 4(u_0, u_1)_{L_g^2} + 2 \cdot 2E(0) \\ & \leq 2 \left\{ 2(u_0, u_1)_{L_g^2} + 2E(0) \right\} \end{aligned}$$

$$(3.14) \leq I_0^2,$$

where

$$(3.15) \quad I_0^2 = 2 \left\{ 2(u_0, u_1)_{L_g^2} + 2E(0) \right\}.$$

Let $\rho = \max\{0, 4\}$, then

$$(3.16) \quad \|u(t)\|_{L_g^2}^2 + \int_0^t \|\nabla u(t)\|^4 dt \leq \rho^{-1} I_0^2.$$

For later use, we introduce the following important function $H(t)$, where

$$(3.17) \quad H(t) = \frac{\|\nabla u_t(t)\|_{L_g^2}^2}{\|\nabla u(t)\|^2} + \|\Delta u(t)\|^2.$$

Next, we multiply equation (1.1) by $-\Delta u_t g$ and integrate over R^N to get (3.18)

$$\begin{aligned} & \int_{R^N} -\Delta u_t u_t g dt + \int_{R^N} \|\nabla u(t)\|^2 \Delta u \Delta u_t dt = 0 \Rightarrow \\ & \frac{d}{dt} \frac{1}{2} \|\nabla u_t(t)\|^2 + \|\nabla u(t)\|^2 \frac{d}{dt} \frac{1}{2} \|\Delta u(t)\|^2 = 0 \Rightarrow \\ & \frac{d}{dt} \|\nabla u_t\|_{L_g^2}^2 + \|\nabla u\|^2 \frac{d}{dt} \|\Delta u\|^2 = 0. \end{aligned}$$

Since we have that $\|\nabla u_0\| > 0$, for $u_0 \neq 0$, we have that $\|\nabla u(t)\| > 0$ near $t = 0$. Let

$$T \equiv \sup\{t \in [0, +\infty) : \|\nabla u(s)\| > 0, 0 \leq s \leq t\}.$$

If $T < +\infty$, we have that $\|\nabla u(T)\| = 0$. We multiply relation (3.19) by $\|\nabla u(t)\|^{-2}$ for $0 \leq t < T$ and we get the following equality

$$(3.19) \quad \frac{d}{dt} H(t) + 2 \left(\frac{(\nabla u, \nabla u_t)}{\|\nabla u\|^2} \right) \frac{\|\nabla u_t\|_{L_g^2}^2}{\|\nabla u\|^2} = 0.$$

Since

$$(3.20) \quad H(0) = \frac{\|\nabla u_1\|_{L_g^2}^2}{\|\nabla u_0\|^2} + \|\Delta u_0\|^2 < 1$$

and

$$(3.21) \quad \frac{|\nabla u(t), \nabla u_t(t)|}{\|\nabla u(t)\|^2} < H(t)^{1/2},$$

we observe that

$$(3.22) \quad \frac{d}{dt} H(t) \leq 0, \quad H(t) \leq H(0),$$

for some $t > 0$, which means that relation (3.22) holds for $0 \leq t < T$, because of contradiction. On the other hand, if $\|\nabla u(T)\| = 0$, we get from (3.22) that $\lim_{t \rightarrow T} \|\nabla u_t(t)\| = 0$. Then, from the uniqueness of the solution (see [20], Proposition 4.1, p. 125) for equation (1.1), we remark that (1.1) has a trivial solution on $[0, T]$, with $\{u(T), u_t(T)\} = \{0, 0\}$. This contradicts the hypothesis that $u_0 \neq 0$. Finally, we conclude that $T = \infty$, that is $\|\nabla u(t)\| > 0$ for $t \geq 0$.

Thus we get, after all these calculations, that equation (1.1) gives a unique local solution u , which belongs to $\bigcap_{k=0}^2 C^k([0, T]; H^{2-k}(R^N))$.

Finally, we have from relations (3.4), (3.17) and (3.22), that

$$\|u(t)\|_{D^{1,2}} + \|u_t(t)\|_{L_g^2} \leq C < \infty \text{ for } t \geq 0.$$

That completes the proof of our Theorem.

References

- [1]. A Arosio and S Garavaldi, *On the mildly degenerate Kirchhoff string*, Math. Methods Appl. Sci., 14(1991), 177–195
- [2]. K J Brown and N M Stavrakakis, *Global Bifurcation Results for a Semilinear Elliptic Equation on all of R^N* , Duke Math. J., 85 (1996), 77–94.
- [3]. H R Crippa, *On local solutions of some mildly degenerate hyperbolic equations*, Nonlinear Analysis TMA, 21 (1993), 565–574.
- [4]. P D'Ancona and Y Shibata, *On global solvability for the degenerate Kirchhoff equation in the analytic category*, Math. Methods Appl. Sci., 17 (1994), 477–489.
- [5]. P D'Ancona and S Spagnolo, *Global solvability for the degenerate Kirchhoff equation with real analytic data*, Invent. Math. 108 (1992), 247–262.

- [6]. P D'Ancona and S Spagnolo, *Nonlinear perturbations of the Kirchhoff equation*, Comm.Pure Appl. Math. 47 (1994), 1005–1029.
- [7]. M Hosoya and Y Yamada, *On some nonlinear wave equations II: global existence and energy decay of solutions*, J. Fac. Sci. Univ. Tokyo Sect. IA Math., 38, (1991), 239–250.
- [8]. Th Gallay and G Raugel, *Scaling Variables and Asymptotic Expansions in Damped Wave Equations*, J Diff Equations, 150 (1998), 42–97.
- [9]. R Ikehata, *Some Remarks on the Wave Equations with Nonlinear Damping and Source Terms*, Nonlinear Analysis TMA, Vol 27, 10, (1996), 1165-1175.
- [10]. N I Karahalios and N M Stavrakakis, *Existence of Global Attractors for semilinear Dissipative Wave Equations on R^N* , J. Differential Equations, 157 (1999), 183–205.
- [11]. N I Karahalios and N M Stavrakakis, *Global Existence and Blow-Up Results for Some Nonlinear Wave Equations on R^N* , Adv. Differential Equations, 6 (2001), 155–174.
- [12]. G Kirchhoff, *Vorlesungen Über Mechanik*, Teubner, Leipzig, 1883
- [13]. T Kobayashi, H Pecher, and Y Shibata, *On a global in time existence theorem of smooth solutions to a nonlinear wave equations with viscosity*, Math. Ann. 296 (1993), 215–234.
- [14]. M Nakao, *Decay of solutions of some nonlinear evolution equations*, J. Math. Anal. Appl., 60 (1977), 542–549.
- [15]. M Nakao, *Energy decay for the quasilinear wave equation with viscosity*, Math. Z., 219, (1995), 289–299.
- [16]. K Nishihara and Y Yamada, *On global solutions of some degenerate quasilinear hyperbolic equations with dissipative terms*, Funkcialaj Ekvacioj, 33 (1990), 151–159.
- [17]. K Nishihara, *Decay properties of solutions of some quasilinear hyperbolic equations with strong damping*, Nonlinear Analysis TMA, 21 (1993), 17–21.
- [18]. P Papadopoulos and N M Stavrakakis, *Compact invariant sets for some quasilinear Nonlocal Kirchhoff Strings on R^N* , Applicable Analysis, 87 (2008), 133–148.
- [19]. R L Pego, *Phase Transitions in one-dimensional Nonlinear Viscoelasticity: Admissibility and Stability*, Arch. Rat. Anal. 97, (1987), 353-394.
- [20]. R Temam, *Infinite-dimensional dynamical systems in mechanics and physics. Applications in Mathematical Science, 2nd Edn*, 68 (1997) (New York: Springer-Verlag).
- [21]. G Todorova, *The Cauchy problem for nonlinear wave equations with nonlinear damping and source terms*, Nonlinear Analysis TMA, 41 (2000), 891–905.
- [22]. E Zeidler, *Nonlinear functional analysis and its applications*, Vol.II, Monotone Operators, Springer-Verlag, 1990.