

# Global Existence, Blow Up And Compact Invariant Sets For Quasilinear Kirchhoff Strings On $R^N$

Perikles Papadopoulos, Niki Lina Matiadou and Georgios Xerogiannakis

Department of Electrical and Electronics Engineering,  
 University of west Attica, Athens, Greece

ppapadop@uniwa.gr, lmatiadou@yahoo.gr, georgiosx@uniwa.gr

**Abstract**—We discuss the asymptotic behavior of solutions for the nonlocal quasilinear hyperbolic problem of Kirchhoff type

$u_{tt} - \phi(x) \|\nabla u(t)\|^2 \Delta u + \delta u_t = |u|^a u$ ,  $x \in R^N$ ,  
 $t \geq 0$ , with initial conditions  $u(x, 0) = u_0(x)$  and  $u_t(x, 0) = u_1(x)$ , in the case where  $N \geq 3, \delta > 0$  and  $(\phi(x))^{-1} = g(x)$  is a positive function lying in  $L^{N/2}(R^N) \cap L^\infty(R^N)$ . It is proved that when the initial energy  $E(u_0, u_1)$  which corresponds to the problem, is non-negative and small, there exists a unique global solution in time in the space  $X_0 =: D(A) \times D^{1,2}(R^N)$ . When the initial energy  $E(u_0, u_1)$  is negative, the solution blows-up in finite time. For the proofs, a combination of the modified potential well method and the concavity method is used. Also, the existence of an absorbing set in the space  $X_1 =: D^{1,2}(R^N) \times L^2_g(R^N)$  is proved and that the dynamical system generated by the problem possess an invariant compact set  $A$  in the same space.

Finally, for the generalized dissipative Kirchhoff's String problem

$$u_{tt} = - \|A^{1/2}u\|_H^2 Au - \delta Au_t + f(u),$$

$$x \in R^N, t \geq 0,$$

with the same hypotheses as above, we study the stability of the trivial solution  $u \equiv 0$ . It is proved that if  $f'(0) > 0$ , then the solution is unstable for the initial Kirchhoff's system, while if  $f'(0) < 0$  the solution is asymptotically stable. In the critical case, where  $f'(0) = 0$ , the stability is studied by means of the central manifold theory. To do this study we go through a transformation of variables similar to the one introduced by R. Pego.

**Keywords**— *Quasilinear Hyperbolic Equations, Global Solution, Blow Up, Dissipation, Potential Well, Cocavity method, Unbounded Domains, Kirchhoff Strings, Generalized Sobolev Spaces, Weighted  $L^p$  spaces.*

## I. Introduction-Preliminaries

We study the following quasilinear hyperbolic initial value problem

$$(1.1) \quad u_{tt} - \phi(x) \|\nabla u(t)\|^2 \Delta u + \delta u_t - |u|^a u = 0,$$

$$(1.2) \quad u(x, 0) = u_0(x), \quad u_t(x, 0) = u_1(x),$$

$x \in R^N, t \geq 0$ , with initial conditions  $u_0, u_1$  in appropriate function spaces,  $N \geq 3$ , and  $\delta \geq 0$ . Throughout the paper we assume that the function  $\phi$  and  $g: R^N \rightarrow R$  satisfy the following condition:

$$(G) \quad \phi(x) > 0, \text{ for all } x \in R^N \text{ and } (\phi(x))^{-1} = g(x) \in L^{N/2}(R^N) \cap L^\infty(R^N).$$

This class will include functions of the form  $\phi(x) \square c_0 + \varepsilon |x|^a, \varepsilon > 0$  and  $a > 0$ , resembling phenomena of slowly varying wave speed around the constant speed  $c_0$ .

G. Kirchhoff in 1883 proposed the so called Kirchhoff string model in the study of oscillations of stretched strings and plates

$$ph \frac{\partial^2 u}{\partial t^2} + \delta \frac{\partial u}{\partial t} = \left\{ p_0 + \frac{Eh}{2L} \int_0^L \left( \frac{\partial u}{\partial x} \right)^2 \right\} \frac{\partial^2 u}{\partial x^2} + f \text{ for } 0$$

$< x < L, t \geq 0$ , where  $u = u(x, t)$  is the lateral displacement at the space coordinate  $x$  and the time  $t$ ,  $E$  the Young modulus,  $p$  the mass density,  $h$  the cross-section area,  $L$  the length,  $p_0$  the initial axial tension,  $\delta$  the resistance modulus and  $f$  the external force (see [10]). When  $p_0 = 0$  the equation is considered to be of *degenerate type*, otherwise it is of *nondegenerate type*.

In the case of bounded domain, T. Kobayashi [11] constructed a unique weak solution by a Faedo-Galerkin method for a quasilinear wave equation with strong dissipation (see also [1, 13]). K. Nishihara [14], has derived a decay estimate from below of the potential of solutions. Also R. Ikehata [4], has shown that for sufficiently small initial data, global existence can be obtained, even when the influence of the source terms is stronger than that of the damping terms. Finally K. Ono [15]

for  $\delta \geq 0$ , has proved global existence and blow up results for a degenerate non-linear wave equation of Kirchhoff type with strong dissipation.

In the case of unbounded domain, P. D'Ancona and S. Spagnolo [2] have shown the global existence of a unique  $C^\infty$  solution for the non-degenerate type with small  $C_0^\infty$  data. N. Karahalios and N. Stavrakakis [5]-[9], have proved global existence and blow-up results for some semilinear wave equations with variable wave speed on all  $R^N$ . T Mizumachi (see [12]), studied the asymptotic behavior of solutions to the Kirchhoff equation with a viscous damping term with no external force. In our previous work (see [16]), we prove global existence and blow-up results of an equation of Kirchhoff type in all of  $R^N$ . Also, in [17] we prove the existence of compact invariant sets for the same equation. Finally, in [18] we study the stability of the trivial solution  $u = 0$  for the generalized Kirchhoff's string equation, using the central manifold theory.

As we will see, the space setting for the initial conditions and the solutions of our problem is the product space  $X_0 =: D(A) \times D^{1,2}(R^N)$ . By

$D^{1,2}(R^N)$  we define the closure of the  $C_0^\infty(R^N)$  functions with respect to the energy norm  $\|u\|_{D^{1,2}} =: \int_{R^N} |\nabla u|^2 dx$ . It is known that

$$D^{1,2}(R^N) = \left\{ u \in L^{\frac{2N}{N-2}}(R^N) : \nabla u \in (L^2(R^N))^N \right\}$$

The weighted Lebesgue space  $L_g^2(R^N)$  is the closure  $C_0^\infty(R^N)$  functions with respect to the inner product  $(u, v)_{L_g^2(R^N)} =: \int_{R^N} guv dx$  (see [3]).

We also have that the operator  $A = -\phi\Delta$  is self-adjoint and therefore graph-closed. Its domain  $D(A)$ , is a Hilbert space with respect to the norm

$$\|Au\|_{L_g^2} = \left\{ \int_{R^N} \phi |\Delta u|^2 dx \right\}^{1/2}. \text{ So, we construct}$$

the following evolution quartet, with compact and dense embeddings:

$$D(A) \subset D^{1,2}(R^N) \subset L_g^2(R^N) \subset D^{-1,2}(R^N).$$

For the positive selfadjoint operator  $A = -\phi\Delta$ , we may define the fractional powers in the following way. For every  $s > 0$ ,  $A^s$  is an unbounded selfadjoint operator in  $L_g^2(R^N)$  with its domain  $D(A^s)$  to be a dense subset in  $L_g^2(R^N)$ . The operator  $A^s$  is strictly positive and injective. Also  $D(A^s)$ , endowed with the scalar product

$$(u, v)_{D(A^s)} = (u, v)_{L_g^2} + (A^s u, A^s v)_{L_g^2},$$

becomes a Hilbert space. We write as usual  $V_{2s} = D(A^s)$  and we have the following

identifications  $D(A^{-1/2}) = D^{-1,2}(R^N)$ ,  $D(A^0) = L_g^2$ ,  $D(A^{1/2}) = D^{1,2}(R^N)$ . Moreover the mapping

$A^{s/2} : V_x \rightarrow V_{x-s}$  is an isomorphism. Furthermore, we

have that the injection  $D(A^{s_1}) \subset D(A^{s_2})$  is compact and dense, for every  $s_1, s_2 \in R, s_1 > s_2$ .

In order to clarify the kind of solutions we are going to obtain for our problem, we give the definition of the weak solution for the problem.

**Definition 1.1** A weak solution of the problem (1.1)-(1.2) is a function  $u$  such that

$$(i) \quad u \in L^2[0, T; D(A)], u_t \in L^2[0, T; D^{1,2}], \\ u_{tt} \in L^2[0, T; L_g^2],$$

(ii) for all  $v \in C_0^\infty([0, T] \times (R^N))$ , satisfies the generalized formula

$$(1.3) \quad \int_0^T (u_{tt}(\tau), v(\tau))_{L_g^2} d\tau \\ + \int_0^T \left( \|\nabla u(\tau)\|^2 \int_{R^N} \nabla u(\tau) \nabla v(\tau) dx \right) d\tau \\ + \delta \int_0^T (u_t(\tau), v(\tau))_{L_g^2} d\tau - \int_0^T f(u(\tau), v(\tau))_{L_g^2} d\tau = 0$$

where  $f(s) = |s|^a s$ , and (iii) satisfies the initial conditions

$$u(x, 0) = u_0(x) \in D(A), u_t(x, 0) = u_1(x) \in D^{1,2}(R^N).$$

In the following section we briefly discuss the results concerning the asymptotic behavior of solutions for the problem (1.1)-(1.2). Among the global existence and blow-up results we also prove existence of a compact functional invariant set. We would like to mention that up to our knowledge, this is the first result concerning existence of functional invariant sets for mathematical models of Kirchoff's strings type.

## II. Global Existence, Blow Up Results and Invariant Sets

In this section we give global existence and blow-up results for the problem (1.1)-(1.2) in the space  $X_0$ .

We also prove existence of an attractor like set. For the proofs we refer on [16], [17]. In order to obtain a local existence result for the problem (1.1)-(1.2), we need information concerning the solvability of the corresponding non-homogeneous linearized problem around the function  $v$ , where

$(v, v_t) \in C(0, T; D(A) \times D^{1,2})$ , is given restricted in the sphere  $B_R$ :

**(2.1)**  
 $u_{tt} - \phi(x) \|\nabla v\|^2 \Delta u + \delta u_t = |v|^a v$ ,  
 $(x, t) \in B_R \times (0, T)$ ,  
 $u(x, 0) = u_0(x), u_t(x, 0) = u_1(x), x \in B_R$ ,  
 $u(x, t) = 0, (x, t) \in \partial B_R \times (0, T)$ ,

$v \in C(0, T; D(A))$  and  $v_t \in C(0, T; \dot{D}^2)$

**Proposition 2.1** Assume that  $u_0 \in D(A), u_1 \in D^{1,2}(R^N)$  and  $0 \leq a \leq 4/(N-2)$ , then the linear wave equation (2.1) has a unique solution such that

$u \in C(0, T; D(A)), u_t \in C(0, T; D^{1,2})$ .

**Proof.** The proof follows the lines of [6, Proposition 3.1]. The Galerkin method is used, based on the information taken from the eigenvalue problem.

Next, we have the following theorem (for the proof we refer to [16]).

**Theorem 2.2** If  $(u_0, u_1) \in D(A) \times D^{1,2}$  and satisfy the non-degenerate condition  $\|\nabla u_0\|^2 > 0$ , then there exists  $T > 0$ , such that the problem (1.1)-(1.2) admits a unique local weak solution  $u$  satisfying:

$u \in C(0, T; D(A)), u_t \in C(0, T; D^{1,2})$ . Moreover, at

least one of the following statements holds true, either

- (i)  $T = +\infty$ , or
- (ii)  $e(u(t)) = \|u_t\|_{D^{1,2}}^2 + \|u\|_{D(A)}^2 \rightarrow \infty$ , as  $t \rightarrow T_-$ .

The next theorem deals with the global existence, blow-up results and the energy decay property of the problem. The proofs of the results are in [16].

First we define as the energy of the problem (1.1)-(1.2) the quantity

$E(t) =: E(u(t), u_t(t)) = \|u(t)\|_{L_g^2}^2 +$

**(2.2)**  $\frac{1}{2} \|u(t)\|_{D^{1,2}}^4 - \frac{2}{a+2} \|u(t)\|_{L_g^{a+2}}^{a+2}$ .

Also, we introduce the potential of the problem (1.1)-(1.2), as

**(2.3)**  $J(u) =: \frac{1}{2} \|u(t)\|_{D^{1,2}}^4 - \frac{2}{a+2} \|u(t)\|_{L_g^{a+2}}^{a+2}$ .

So, we get the following relation

**(2.4)**  $E(t) = \|u_t(t)\|_{L_g^2}^2 + J(u)$ .

Finally, we introduce a modified version of the modified potential well used in [6] (see also [13]), by

**(2.5)**

$W =: \left\{ u \in D(A); K(u) = \|u\|_{D^{1,2}}^4 - \|u\|_{L_g^{a+2}}^{a+2} > 0 \right\} \cup \{0\}$ .

**Theorem 2.3** Assume that  $N = 3, 8/3 < a < 4, u_0 \in W(\subset D(A))$  and  $u_1 \in D^{1,2}$ .

Also suppose that the following inequality holds

**(2.6)**  $E(u_0, u_1) \leq \left( \frac{1}{C_0 \mu_0^{p_1}} \right)^{1/p_2}$ , if  $8/3 < a < 4$  and

$p_2 > 0$ . Then **a)** for  $p_1 =: \frac{2(a+2)-3a}{2}$  and

$p_2 =: \frac{3a-8}{8}$ , there exists a unique global solution

$u \in W$  of the problem (1.1)-(1.2) satisfying

$u \in C([0, +\infty); D(A))$  and  $u_t \in C([0, +\infty); D^{1,2})$ .

**b)** Moreover, this solution obeys the following energy estimates

$\|u_t\|_{L_g^2}^2 + d_*^{-1} \|\nabla u\|^4 \leq E(u, u_t) \leq$

**(2.7)**  $\left\{ E(u_0, u_1)^{-1/2} + d_0^{-1} [t-1]^+ \right\}^{-2}$ ,

where  $d_* = \frac{2(a+2)}{a-2}$  and  $d_0 \geq 1$ , that is

**(2.8)**  $\|\nabla u\|^4 \leq C_*(1+t)^{-1}$ ,

where  $C_*$  is some constant depending on  $\|u_0\|_{D^{1,2}}^4$  and  $\|u_1\|_{L_g^2}$ .

**c)** Suppose that  $a \geq 2, N \geq 3$  and the initial energy  $E(u_0, u_1)$  is negative. Then there exists a time  $T$ , where

$0 < T \leq a^{-2} (-E(u_0, u_1))^{-1}$

**(2.9)**  $\left[ \left\{ \left( 2\delta \|u_0\|_{L_g^2}^2 - a(u_0, u_1)_{L_g^2} \right)^2 \right. \right.$

$\left. \left. + a^2 (-E(u_0, u_1)) \|u_0\|_{L_g^2}^2 \right\}^{1/2} + \right.$

$\left. 2\delta \|u_0\|_{L_g^2}^2 - a(u_0, u_1)_{L_g^2} \right]$ ,

such that the (unique) solution of the problem (1.1)-(1.2) blows-up at  $T$ , i.e.,

**(2.10)**  $\lim_{t \rightarrow T_-} \|u(t)\|_{L_g^2}^2 = +\infty$ .

The existence of an absorbing set in  $X_0$  is given below. The proofs of the results are in the work [17].

**Lemma 2.4** Assume that  $\rho_1 > 4a^{-1/2} R^2 c_3^2, 0 \leq a < 2/(N-2), N \geq 3$  and  $\|\nabla u_0\| > 0$ . Then the unique local solution defined by Theorem 2.1 exists globally in time.

**Remark 2.5** (Global Solutions) From the last Lemma 2.4, we may observe that solutions of the problem

(1.1)-(1.2), (given by Theorem 2.2), belong to the space  $C_b(R_+, X_0)$ , i.e., we have achieved global solutions for the given problem. Let us remark that, in the Theorem 2.3, using a modified potential well technique, we have proved global existence results under the conditions  $N=3, 8/3 < a < 4$  and the initial energy  $E(0)$  been non-negative and small. On the other hand, in Lemma 2.4, we could achieve global results for different type of nonlinearities, i.e.,  $a \in (0, 2/(N-2))$ , but for any  $N \geq 3$  and independently of the sign of the initial energy  $E(0)$ .

Lemma 2.4 has an immediate consequence:

**Remark 2.6** A nonlinear semigroup  $S(t): X_0 \rightarrow X_0, t \geq 0$ , may be associated to the problem (1.1)-(1.2) such that for  $\psi = \{u_0, u_1\} \in X_0, S(t)\psi = \{u(t), u_t(t)\}$  is the weak solution of the problem (1.1)-(1.2). Moreover the ball  $B_0 =: B_{X_0}(0, \bar{R}_*)$  for any  $\bar{R}_* > R_*$ , where  $R_*$  is defined by Lemma 2.4, is an **absorbing set** for the semigroup  $S(t)$  in the energy space  $X_0 \subset X_1$ , compactly.

In the rest of the paper we show that the  $\omega$ -limit set of the absorbing set  $B_0$  is a compact invariant set. To this end, we need to decompose the semigroup  $S(t)$ , in the form  $S(t) = S_1(t) + S_2(t)$ , where for a suitable bounded set  $B \subset X_0$ , the semigroups  $S_1(t), S_2(t)$  satisfy the following properties:

**(S1)**  $S_1(t)$  is uniformly compact for  $t$  large, i.e.,  $\cup_{t \geq t_0} S_1(t)B$  is relatively compact in  $X_1$ .

**(S2)**  $\sup_{k \in B} \|S_2(t)k\|_{X_1} \rightarrow 0$ , as  $t \rightarrow \infty$ .

As a consequence of the above properties we have the following result

**Theorem 2.7** Let  $\phi$  satisfy hypothesis (G). Then the semigroup  $S(t)$  associated with the problem (1.1)-(1.2) possesses a functional invariant set  $A = \omega(B_0)$ , which is compact in the weak topology of  $X_1$ .

**Remark 2.8** We have that  $X_0$  is compactly embedded in  $X_1$ , so the set  $\cup_{t \geq t_0} S_1(t)B$  is compact with respect to the strong topology in  $X_1$ . For the functional invariant compact set  $A = \omega(B_0)$ , we observe that  $(u_0, u_1) \in A$ , if  $|\nabla u_0| > 0$ . So,  $A$  is an **attractor like set**.

Finally, in the following section we study the stability of the initial solution  $u=0$  for the generalized Kirchhoff equation.

### III. Stability Results

We consider the generalized quasilinear dissipative Kirchhoff's String problem

$$u_{tt} = -\|A^{1/2}u\|_H^2 Au - \delta Au_t + f(u),$$

$$x \in R^N, t \geq 0,$$

under the same initial conditions as above and  $H$  is a Hilbert space. First, we prove existence of solution for our problem, under small initial data (for the proof we refer to [18]).

**Theorem 3.1.** (Local Existence) Let  $f(u)$  a  $C^1$ -function such that  $|f(u)| \leq k_1 |u|^{a+1}, |f'(u)| \leq k_2 |u|^a, 0 \leq a \leq 4/(N-2), N \geq 3$ .

Consider that  $(u_0, u_1) \in D(A) \times V$  and satisfy the non-degenerate condition

$$(3.1) \quad \|A^{1/2}u_0\| > 0.$$

Then there exists  $T_0 > 0$  such that our problem admits a unique local weak solution  $u$  satisfying  $u \in C(0, T; D(A))$  and  $u_t \in C(0, T; V)$ .

The linearized equation of the system around the solution  $u=0$  is

$$(3.2) \quad \bar{u}_t + A^* \bar{u} = 0,$$

where

$$(3.3) \quad \bar{u}_t = (w, v)^T \text{ and } A^* = \begin{bmatrix} \delta A & -f'(0) \\ -1 & 0 \end{bmatrix}.$$

So, in order to study the stability of the solution, we study the spectrum of the operator  $A^*$ . The characteristic polynomial of  $A^*$  is

$$\begin{vmatrix} -\delta \lambda_j + \mu_j & f'(0) \\ 1 & \mu_j \end{vmatrix} = 0,$$

or equivalently

$$\mu_j^2 - \delta \lambda_j \mu_j - f'(0) = 0.$$

Let,  $\Delta = \delta^2 \lambda_j^2 + 4f'(0)$ . Then according to the sign of  $f'(0)$ , we have the following cases:

**I)** Let  $f'(0) > 0$ , then we have that 0 is unstable for the initial Kirchhoff's system.

**II)** Let  $f'(0) < 0$ . This implies that the operator  $A^*$  admits two real eigenvalues, which are both positive. Thus we obtain that the solution  $u=0$

is asymptotically stable for the initial Kirchhoff's system.

**III)** Let  $f'(0) = 0$ . In this case we use the central manifold theory in order to study the stability of the initial solution  $u = 0$ . Making use of the change of variables similar to what is found by Pego (see [20]), namely

$$(3.4) \begin{cases} p(x, t) = A^{-1/2} u_t, \\ q(x, t) = -\delta A^{1/2} u - p, \end{cases}$$

we can rewrite (3.2)-(3.3) in the form of a reaction-diffusion system:

$$(3.5) \begin{cases} p_t(x, t) = -\delta A p + \left(\frac{1}{\delta^3} \|p + q\|_H^2\right)(p + q) + A^{-1/2} f(u), \\ q_t(x, t) = -\left(\frac{1}{\delta^3} \|p + q\|_H^2\right)(p + q) - A^{-1/2} f(u), \\ p(x, t) = 0, t > 0, \\ p(x, 0) = p_0(x), q(x, 0) = q_0(x), \end{cases}$$

where  $p + q = -\delta A^{1/2} u$ .

In order to prove the existence of a local central manifold we need the following result:

**Proposition 3.2.** For some neighbourhood  $U$  of 0 in  $X^{1/2} =: V \times H$ , system (3.5) has a local central manifold defined by

$$W_{loc}^c(0) = \{ \xi + \eta \mid \xi = h^c(\eta), \xi \in X_+^{1/2} \cap U, \eta \in X_0 \cap U \}$$

where we have that  $h^c(0) = Dh^c(0) = 0$ .

We get that the central manifold is approximated in the following form

$$(3.6) \begin{cases} h^c(q) = \frac{1}{\delta^4} \|q\|_H^2 A^{-1} q + \\ \frac{2A^{-3/2} f(u)}{\delta} + O(\|q\|_H^4). \end{cases}$$

Solutions on the central manifold satisfy

$$(3.7) \begin{cases} p(t) = h^c(q(t)), \\ q_t(t) = -\frac{1}{\delta^3} \|h^c(q) + q\|_H^2 (h^c(q) + q). \end{cases}$$

From system (3.7), we obtain that the stability of the solution  $u = 0$  depends on  $f$ . Thus we have the following cases:

**(i):** if  $f(u_0) < 0$ , then we get that  $(p, q) = (0, 0)$  is unstable, so  $u = 0$  is also unstable for the initial Kirchhoff's system,

**(ii):** if  $f(u_0) > 0$ , then  $(p, q) = (0, 0)$  is Asymptotically stable, so  $u = 0$  is also asymptotically stable for the initial system,

**(iii):** if  $f(u_0) = 0$ , we have that solutions on the central manifold satisfy the following system

$$\begin{cases} p(t) = h^c(q(t)), \\ q_t(t) = -\frac{1}{\delta^3} \|q\|_H^2 q + O(\|q\|_H^5). \end{cases}$$

So, we obtain that  $(p, q) = (0, 0)$  is stable, that is,  $u = 0$  is also stable for the initial Kirchhoff's system.

## References

- [1]. P D'Ancona and Y Shibata, *On global solvability for the degenerate Kirchhoff equation in the analytic category*, Math. Methods Appl. Sci., 17 (1994), 477-489.
- [2]. P D'Ancona and S Spagnolo, *Nonlinear perturbations of the Kirchhoff equation*, Comm.Pure Appl. Math. 47 (1994), 1005-1029.
- [3]. K J Brown and N M Stavrakakis, *Global Bifurcation Results for a Semilinear Elliptic Equation on all of  $R^N$* , Duke Math. J., 85 (1996), 77-94.
- [4]. R Ikehata, *Some Remarks on the Wave Equations with Nonlinear Damping and Source Terms*, Nonlinear Analysis TMA, Vol 27, 10, (1996), 1165-1175.
- [5]. N I Karachalios and N M Stavrakakis, *Existence of Global Attractors for semilinear Dissipative Wave Equations on  $R^N$* , J. Differential Equations, 157 (1999), 183-205.
- [6]. N I Karachalios and N M Stavrakakis, *Global Existence and Blow-Up Results for Some Nonlinear Wave Equations on  $R^N$* , Adv. Differential Equations, 6 (2001), 155-174.
- [7]. N I Karachalios and N M Stavrakakis, *Asymptotic Behavior of Solutions of Some Nonlinearly Damped Wave Equations on  $R^N$* , Topological Methods in Nonlinear Analysis, Vol 18, (2001), 73-87.
- [8]. N I Karachalios and N M Stavrakakis, *Estimates on the Dimension of a Global Attractor for a Semilinear Dissipative Wave Equation on  $R^N$* , Discrete and Continuous Dynamical Systems, Vol 8, (2002), 939-951.

- [9]. N I Karachalios and N M Stavrakakis, *Global Attractor for the Weakly Damped Driven Schrodinger Equation in  $H^2(R)$* , NoDEA, (2002), 347-360.
- [10]. G Kirchhoff, *Vorlesungen Uber Mechanik*, Teubner, Leipzig, 1883.
- [11]. T Kobayashi, H Pecher, and Y Shibata, *On a global in time existence theorem of smooth solutions to a nonlinear wave equations with viscosity*, Math. Ann. 296, (1993), 215-234.
- [12]. T Mizumachi, *The Asymptotic Behavior of Solutions to the Kirchhoff Equation with a Viscous Damping Term*, Journal of Dynamics and Differential Equations, 9, (1997), 211-247.
- [13]. M Nakao, *Energy decay for the quasilinear wave equation with viscosity*, Math. Z., 219, (1995), 289-299.
- [14]. K Nishihara, *Decay properties of solutions of some quasilinear hyperbolic equations with strong damping*, Nonlinear Analysis TMA, 21, (1993), 17-21.
- [15]. K Ono, *On global existence, asymptotic stability and blowing up of solutions for some degenerate non-linear wave equations of Kirchhoff type with a strong dissipation*, Math. Meth. Appl. Sci., 20, (1997), 151-177.
- [16.] P G Papadopoulos and N M Stavrakakis, *Global Existence and Blow-Up Results for an Equation of Kirchhoff Type on  $R^N$* , Topological Methods in Nonlinear Analysis, 17, (2001), 91-109.
- [17]. P G Papadopoulos and N M Stavrakakis, *Compact Invariant Sets for Some Quasilinear Nonlocal Kirchhoff Strings on  $R^N$* , Applicable Analysis, 87 (2008), 133–148.
- [18]. P G Papadopoulos and N M Stavrakakis, *Central Manifold Theory for the Generalized Equation of Kirchhoff Strings on  $IRN$* , Nonlinear Analysis TMA, 2005, Vol 61, pp 1343-1362.
- [20]. P G Papadopoulos and N M Stavrakakis, *Strong Global Attractor for a quasilinear nonlocal wave equation on  $IRN$* , Electronically Journal of Differential Equations (EJDE), 2006, Vol 77, pp 1-10.
- [21]. R L Pego, *Phase Transitions in one-dimensional Nonlinear Viscoelasticity: Admissibility and Stability*, Arch. Rat. Anal. 97, (1987), 353-394.