

Complex Formation in a Liquid-Liquid Extraction System Containing Nickel (II), 2,6-Dithiol-4-methylphenol, and Diphenylquandine

Kuliev Kerim Avaz¹, Mamedova Rena Almaz¹, Ismailova Shahnaz Yusif²

1-Department of Chemistry, Azerbaijan State Pedagogical University

U. Gadjibekov street 68. Baku, AZ 1000, Azerbaijan

2-Azerbaijan Medical University

kerim.kuliev.69@mail.ru

Abstract—Complex formation and liquid-liquid extraction were studied in a system containing nickel(II), 2,6-Dithiol-4-methylphenol (DTMP), Diphenylquandine (DPG), water, and chloroform. The effect of some experimental parameters (pH, shaking time, concentration of DTMP, and concentration of DPG) was systematically investigated, and the optimum conditions for cobalt extraction as an ion-association complex, $(DPGH)_2[Ni(DTMP)_2]$, were found. The following key equilibrium constants were calculated: constant of association ($\log K = 9.96 \pm 0.05$), constant of distribution ($\log K_D = 2.09 \pm 0.01$), and constant of extraction ($\log K_{ex} = 12.05 \pm 0.09$). Beer's law was obeyed for Ni concentrations up to $3.2 \mu\text{g} \times \text{mL}^{-1}$ with a molar absorptivity of $3.24 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$ at $\lambda_{max} = 520 \text{ nm}$. Some additional characteristics, such as limit of detection, limit of quantification, and Sandell's sensitivity, were estimated as well.

Keywords—Spectrophotometry; Nickel (II)-DTMP complex and different samples

1. INTRODUCTION

Nickel is widely used in electroplating, the manufacture of Ni-Cd batteries, rods for arc welding, pigments of paints, ceramic, surgical and dental prostheses, magnetic tapes and computer components and nickel catalysts. Nickel enters waters from dissolution of industrial processes and waste disposal. Nickel is essential constituent in plant urease. Jack beans and soybeans generally contain high concentration of nickel [1].

Nickel-containing sewage is harmful after ingress into water. This fact explained the importance of the monitoring of nickel concentration in natural and waste water samples. Flame and graphite furnace atomic absorption spectrometry and spectrophotometric methods provides accurate and rapid determination of nickel in natural and waste waters [2]. However, very frequently a direct determination cannot be applied due to low concentration of analyte or matrix interferences.

Several analytical methods have been used for nickel determination, including flame atomic absorption spectrophotometry [3], graphite furnace atomic absorption spectrometry [4], electrothermal atomic absorption spectrometry [5], atomic fluorescence spectrometry [6], inductively coupled plasma-optical emission spectrometry [7], and spectrophotometry [8.;

9. 10.11.12].

The methods involving spectrophotometry are rather popular due to their simplicity, inexpensive instrumentation and easy automation. Nickel (II) chelates of several ligands like diphenylcarbazone, diphenylthiocarbazone, 8-

quinolinol and its substituted analogues produce intense absorption in the optical spectra, in the visible range, which is characteristic of the ligand itself. Structural changes occurred in the nickel (II) chelates on addition of nitrogen bases forms adduct which gives rise to profound spectral changes. Such bathochromic and hypsochromic shifts observed in the visible region of the spectra of nickel (II) chelates of dithizone, dipheylcarbazone, 8-quinolinol, methylsubstituted 8-quinolinols, substituted diphenylcarbazones and dithizones on addition of nitrogen bases. These were employed for the determination of adduct formation constants [13,14,15].

Oxyphenolate and dithiophenolate complexes of cobalt are insoluble in chloroform, while mixed-ligand complexes with hydrophobic amines and aminophenols easily dissolve in various organic solvents [16,17,18,19,20,21,22, 23].

In the present paper we study the complex formation in a liquid-liquid extraction system containing Ni(II), 2,6-dithiol-4-methylphenol (DTMP) and diphenylguanidine (DPG), and show the potential of this system for Ni(II) determination in real samples. HITP, an iodine-containing analogue of the known analytical reagents 2,6-dithiolphenol and 2,6-dithiol-4-*tert*-butylphenol, has -OH and -SH groups in orthoposition. Hence, we expected that this reagent would form stable chelate structures with metal ions, like Ni(II), which have ionic radii close to 0.07-0.08 nm.

2. Experimental

2.1. Reagents and Apparatus

The standard solution (1mg / ml) Ni (II) was prepared by dissolving in water an exact linkage $(\text{NH}_4)_2\text{Ni}(\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ in water containing 2 ml conc. H_2SO_4 [15].

Solutions of DTMP and DPG in chloroform (0.01M) were used. DTMP were synthesized according to the procedure [24]. Their purity was verified by melting point determination and paper chromatography. To create the optimal acidity, 0.1M solutions of KOH and HCl or ammonium acetate buffers were applied. Acetate buffer solution, prepared by mixing of $2 \text{ mol} \times \text{L}^{-1}$ aqueous solutions of CH_3COOH and NH_4OH .

The extractant was purified chloroform.

The absorbance of the extracts was measured using a SF 26 spectrophotometer (USSR) and KFK 2 photocolimeter (USSR). Glass cells with optical path of 5 or 10 mm were used. pH of aqueous phase was measured using an I-120.2 potentiometer with a glass electrode. Muffle furnace was used for dissolution of the samples.

2.2. General Procedure

2.2.1. General Procedure for the Determination of Nickel (II)

Portions of stock solutions of Nickel (II) varying from 0.1 to 1.0 mL with a 0.1-mL step, a 2.2 mL portion of a 0.01 M solution of DTMP, and a 2.5 mL portion of a 0.01M solution of Am were placed in to calibrated test tubes with ground-glass stoppers (the volume of the organic

phase was 5 mL). The required value of pH was adjusted by adding 1M HCl. The volume of the aqueous phase was increased to 20 mL using distilled water. In 15 minnute after the complete separation of the phases, the organic phase was separated from the aqueous phase and the absorbance of the extracts was measured on KFK-2 at room temperature and 440 nm ($l=0.5\text{cm}$).

2.2.2. Determination of Nikel (II) in steel

A weighed sample of 0.2 g was dissolved in 20 ml of H_2SO_4 (1: 1) was oxidized with a few drops of concentrated nitric acid and evaporated twice lo vapor SO_3 . The precipitated salt was dissolved in 20 ml of 15% tartaric acid under heating, the solution was cooled, adjusted with water to 100 ml in a volumetric flask, stirred and filtered. An aliquot of 5 ml was put into a separatory funnel, was added 1 ml of 10% hydroxylamine solution, 1 ml of 3% ascorbic acid and was determined cobaltusing the proposed procedures.

2.3.3. Determination of Ni (II) in sewage water and Bottom sediments

Il taken for analysis of waste water is evaporated to obtain a precipitate, do not boil. The precipitate was dissolved in 5 ml of HNO_3 , was transferred to a 50 ml flask and diluted to the mark with water.

3. Results and Discussion

Ni(II) reacts with 2,6-dithiol-4-methylphenols (DTMP) and gives a yellow colored complexes. These complexes are

insoluble in non-polar solvents. Experiments on electromigration in a U-shaped tube and on sorption on EDE-10P (EDE- ethylenediamine, epichlorohydrin; 10- serial number of the brand: P- means that the matrix has a macroporous structure) anion exchangers have demonstrated the anionic nature of single-ligand complexes, in the electromigration study of the complexes, it was found that the yellow dithiophenolate complexes of Nikel (II) moved to the cathode. When the sign of the charge of the single-ligand complexes was determined by ion chromatography, the EDE-10P anion exchanger completely absorbed the colored component of the solution. When hidrophob amins (Am) were introduced into the system, the extraction of these compounds into the organic phase as a mixed-ligand complex (MLC) was observed.

3.1. The Choice of the Extractant

For the extraction of complexes we used CHCl_3 , CCl_4 , C_6H_6 , $\text{C}_6\text{H}_5\text{CH}_3$, $\text{C}_6\text{H}_4(\text{CH}_3)_2$, $\text{C}_2\text{H}_4\text{Cl}_2$, isobutanol and isopentanol. The extractivity of the complexes was estimated by the distribution coefficient and recovery. Chloroform, dichloroethane and chlorobenzene appeared to be the best extractants. All the further investigations were carried out with chloroform. The concentration of nikel in the organic phase was determined with Dimethylglyoxime [15] by photometric measurements after back extraction, while in the aqueous phase it was determined by the difference. The basicity of Am hardly influences the recovery of niobium. After a single

extraction with chloroform, 96.9-99.2% of cobalt was extracted as an ion associate.

3.2. Absorption Spectra

Spectra of the extracted ternary Ni-DTMP-DPG complex and the blank are shown in Figure 2. A maximum is recorded at 520 nm, where the blank absorbs insignificantly. It is shifted to 10 nm as compared to the maximum of the binary Ni-PAR chelate existing in aqueous medium (in the pH interval from 3.2 to 9.5): 510 nm [9, 27, 29, 31, 39]. The observed bathochromic effect is small and gives us grounds to suggest the formation of a ternary compound of the ion-association type.




Fig.2. Absorption of mixed-ligand complexes
 $C_{Ni}=3.44 \times 10^{-5}$ M. $c_{DTMP} = c_{Am}=1.0 \times 10^{-3}$ M,
 KFK-2, 540 nm, $l=1.0$ cm.

3.2. Effect of pH

Results showed that the optimal pH for the extraction of Co with PAR and Nt is 6.5–7.5 (Figure 3). A buffer solution with a concentration of 2 mol L^{-1} was applied to control pH. The use of 0.25–5 mL of the buffer solution per 10 mL (final aqueous solution) was found to

give a constant absorbance. All further experiments were carried out with 5 mL buffer solution with $\text{pH} = 6.5\text{--}6.6$.




Fig. 2. Absorbance of mixed-ligand complexes as a function of the pH of the aqueous phase.
 $C_{Ni}=2.035 \cdot 10^{-5}$ M. $c_{DTMP} = c_{Am}=1.0 \times 10^{-3}$ M, 540 nm,
 $l=0.5$ cm.

3.3. Effect of Reagents' Concentrations

For the formation and extraction of MLC, a 20-25-fold excess of complexing reagents is required; for example, the optimal conditions for formation and extraction of these compounds are provided by 1.0×10^{-3} M DTMP and $(0.92) \times 10^{-3}$ M DPG. A large excess of hydrophob amin interferers with the determination. However it was found that the presence of excess of the reagent solution does not alter the absorbance of the color reaction.

3.4. Effect of Shaking Time

The extraction equilibrium is reached for a short shaking time (about 5 seconds). It was found that a shaking time longer than 1 min can

bring about to a slight decrease (5-6%) of the absorbance values. To avoid this disadvantage and to guarantee complete transfer of the complex into organic phase, even under nonoptimum conditions, the authors extracted in their experiments for 15–20 seconds. Unlike single-ligand complexes, mixed-ligand complexes of Ni(II) with DP and Am were stable in aqueous and organic solvents and did not decompose for two days, or over a month after extraction. The required duration of the phase contact was 10 min.

3.5. Stoichiometry of the Complexes and the Mechanism of Complexation

The molar DTMP-to-Ni(II) and DPG-to-Ni(II) ratios were determined by the mobile equilibrium method [25], molar ratio method [25], and the method of Asmus [25]. The results showed that the ternary complex has a composition of 1 : 2 : 2: (DPGH)₂[Ni(DTMP)₂]. In this formula, DTMP is in deprotonated form (DTMP²⁻), while DPG is in protonated form (DPGH⁺). The mentioned formula and the known properties of DTMP [45–47] and DPG [48] fit well to the observed pH curve of the ternary complex presented in Figure 3: (1) at pH values lower than pH_{opt} DTMP is hardly possible to be in its DTMP²⁻ form; (2) at pH values higher than pH_{opt} DPG is hardly possible to be in its DPGH⁺ form.




Fig. 3. Determination of the ratio of components by method of Asmus for

1. Ni(II):DTMP; 2. Ni(II) : DPG

$C_{\text{Ni(II)}} = 3.44 \cdot 10^{-5} \text{ mol L}^{-1}$, $C_{\text{Am}} = C_{\text{DTMP}} = 1.0 \cdot 10^{-3} \text{ mol} \cdot \text{L}^{-1}$, pH = 6, 540 nm, l=0.5 cm

The disappearance of the pronounced absorption bands in the 3200–3600 cm^{-1} with a maximum at 3460 cm^{-1} observed in the spectrum of DTMP, says that the -OH group is involved in the formation of the complex. The observed decrease in the intensity, absorption bands in the area 2580 cm^{-1} shows that one of the -SH groups involved in the formation of coordination bond in the ionized state. Detection of the absorption bands at 2380 cm^{-1} indicates the presence of a protonated diphenylguanidine [26,27].




Fig. 4. IR spectrum of the complex Ni(II)-DTMP-DPG

Structure extractable complexes can be represented as in Figure 5.





Fig. 5. Structure of complex $[\text{Ni}(\text{DTMP})_2](\text{DPGH})_2$


Calculation of extent of polymerization of complexes was carried out on the equation [28]. The made calculations showed that MLC in an organic phase won't be polymerized and are in a monomeric form ($\nu=1,05-1,12$). It was found using the Nazarenko method that Ni(II) in the complexes was present in the form of Ni^{2+} . The number of protons replaced by cobalt in one DP molecule appeared to be one [29, 30].

Several equilibrium processes should be taken into account for the system of $[\text{Ni}(\text{DTMP})_2]^{2-}$, DPGH^+ , water, and chloroform.


Formation of ion-association complex in the aqueous phase:



Distribution of the complex between the aqueous and the organic phase:



Extraction from water into chloroform:



Based on these equation:

$$\beta = \frac{\{[\text{Ni}(\text{DTMP})_2](\text{DPGH})_2\}_{\text{org}}}{\{[\text{Ni}(\text{DTMP})_2]^{2-}\}_{\text{aq}}\{[\text{DPGH}^+]^2\}_{\text{aq}}} \quad (4)$$

$$K_D = \frac{\{[\text{Ni}(\text{DTMP})_2](\text{DPGH})_2\}_{\text{org}}}{\{[\text{Ni}(\text{DTMP})_2]^{2-}\}_{\text{aq}}\{[\text{DPGH}^+]^2\}_{\text{aq}}} \quad (5)$$

$$K_{\text{ex}} = \frac{\{[\text{Ni}(\text{DTMP})_2](\text{DPGH})_2\}_{\text{org}}}{\{[\text{Ni}(\text{DTMP})_2]^{2-}\}_{\text{aq}}\{[\text{DPGH}^+]^2\}_{\text{aq}}} \quad (6)$$

The association constant β was determined by several independent methods: Holme-Langmihir method [31], Harvey-Manning method [32], and mobile equilibrium method [33]. The distribution constant K_D was calculated from the absorption values obtained after single and double extraction as described above. The extraction constant K_{ex} was calculated by the formula $K_{\text{ex}} = K_D \times \beta$. The recovery factor R was estimated by the dependence $R\% = 100K_D/(K_D + 1)$ and the following value was obtained $R = 99.2\%$. All experiments were performed at room temperature of $\sim 22^\circ\text{C}$ and the calculations were carried out at a probability of 95%.

3.6. Influence of Interfering Ions

The effect of various ions and reagents on the extraction-spectrophotometric determination of 5 mg nickel (II) is summarised in Table 1. It can be assumed that large amounts of alkaline ions, alkaline-earth ions, NH_4^+ , Cl^- , $\text{S}_2\text{O}_3^{2-}$, F^- , NO_3^- , SO_4^{2-} , ClO_4^- , PO_4^{3-} , tartrate, citrate, oxalate and tiron; moderate amounts of Cr(VI), Cr(III), Zn(II) and Cd(II); and small amounts of Mn(II), Sn(II), Cu(II), Al(III), ascorbic acid and SCN^- are tolerable. Co (II), Fe(II,III), V(IV,V), W(VI), Mo(VI), Ti(IV) and Nb(V) interfere determination of Ni(II). However, the interfering effect of some of these ions can be reduced by masking with oxalate, citrate or EDTA (see Table 3).

Table 1. Influence of interfering ions on the determination of nickel(II) as MLC with DTMP and DPG (30,0 μg Ni added)

Ion	Molar excess of the ion	Masking agent	Found Ni, μg	S_r
Co(II)	50		30,2	2
Ni(II)	50		29,8	4
Fe(II)	200		29,8	3
Cd(II)	200		30,3	4
Al(III)	180		29,6	3
Fe(III)	60	Ascorbic acid	30,2	5
Zr(IV)	50		29,8	3
Cu(II)	25	Thiourea	30,1	5
Hg(II)	40		30,2	5
Ti(IV)	30	Ascorbic acid	29,8	4
V(IV)	20		29,6	3
Mo(VI)	10	EDTA	30,4	4
W(VI)	50		29,8	4
Cr(III)	120		29,8	4
Nb(V)	50	$C_2O_4^{2-}$	29,8	4
Ta(V)	50	$C_2O_4^{2-}$	29,7	5
UO_2^{2+}	50		30,2	4
Salicylic acid	25		30,4	4
Sulphosalicylic acid	30		29,8	5
Ascorbic acid	120		30,0	3
Tartaric acid	120		30,2	6
Oxalate	40		30,7	4
Fluoride	35		29,6	7
Phosphoric acid	30		30,3	3
Thiourea	20		31,2	5

3.7. Beer's Law, Molar Absorptivity, and other Analytical Characteristics.

The range of adherence to Beer's law was studied at the optimum conditions (Table 1). The linearity is observed up to $3.2 \mu\text{g ml}^{-1}$ of Ni with a correlation coefficient of 0.9995. The obtained straight line equation is $Y = 0.045 + 0.257 X$. The molar absorptivity was calculated to be $3.24 \times 10^4 \text{ L mol}^{-1} \text{ cm}^{-1}$. This value could compete successfully with the ones obtained for similar DTMP-containing complexes (Table 3). The limit of detection (LOD) and limit of quantification (LOQ) were estimated at 3 times and 10 times standard deviation of the intercept divided by the slope. Sandell's sensitivity was

calculated as well. The values of the above-mentioned characteristics are included in Table 2. Sandell's sensitivity ($\mu\text{g} \cdot \text{cm}^{-2}$) 1.73. [34].




Fig. 6. Analytical determination of Ni (II);

$C_{DTMP} = 1 \times 10^{-3} \text{ M}$; $C_{Am} = 0.92 \times 10^{-3} \text{ M}$; $\text{pH} = 3$;

$\lambda = 540 \text{ nm}$, $l = 0.5 \text{ cm}$.

Table 2. Optical characteristics, precision and accuracy of the spectrophotometric determination of Ni(V) with DTMP and DPG

Parameter	Value
Color	red
The pH range of education and extraction	2,5-9,5
The pH range of maximum extraction	5,5-6,5
Concentration of DTMP: $\text{mol} \cdot \text{L}^{-1}$	1.0×10^{-3}
Concentration of DPG: $\text{mol} \cdot \text{L}^{-1}$	0.92×10^{-4}
Organic solvent	Chloroform
Extraction time	5 min
λ_{max} (nm)	520
Molar absorptivity ($\text{L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$)	$3.24 \cdot 10^4$
Sandell's sensitivity ($\mu\text{g} \cdot \text{cm}^{-2}$)	0.00173
R, %	99.2
The equation of calibration curves	$0.045 + 0.257$
$\lg K_e$	7.85
$\lg K_{\text{Ex}}$	12.05 ± 0.09
Stability constant (β)	9.96 ± 0.05
Beer's law range ($\mu\text{g} \cdot \text{ml}^{-1}$)	0.05-3.2
Correlation coefficient	0.9964
Limit of detection (LOD): $\text{ng} \cdot \text{mL}^{-1}$	12
Limit of quantification (LOQ): $\text{ng} \cdot \text{mL}^{-1}$	40

The proposed method compares favourably with the existing ones (Table 3) and offers the advantages of better simplicity, rapidity, sensitivity and selectivity.

Table 3. Comparative characteristics of the procedures for determining nikel

Reagent	pH (solvent)	λ , nm	$\epsilon \cdot 10^{-4}$	Beer's law range ($\mu\text{g} \cdot \text{ml}^{-1}$)
Dimethylglyoxime [35]	12	470		0.26-2.1
N-ethyl-3-carbazolecarboxaldehyde-3-thiosemicarbazone.[36]	6.0	400	1.114	
7-Methyl-2-chloroquinoline-3-carbaldehyde thiosemicarbazone [37]	6.0	410	1.67	
Thiazole-2-carbaldehyde 2-Quinolyldrazone [38]	8.7 -9.5	522	7.17	0-0.7
pyridoxal-4-phenyl-3-thiosemicarbazone [39]	4-6	430	1.92	0.5-5
4-gidroksiben'aldegid-4-brom fenilgidrazin [40]	4	497	12.85	0.01-0.1
DTMP+DPG	6.5-7.5	527	3.24	0.05-3.2

3.8. Analytical Applications

The proposed method under the already established optimum conditions was applied for the determination of Ni(II) in various objects. The results presented in Table 4 and Table 5 indicate the successful applicability of the proposed method to real sample analysis.

Table 4. Determination of Nikel in steel ($n=5$, $P=0.95$)

Reagent	\bar{X} , %	S_x	ϵ	S_r	μ
$8X\Phi (C16^6)$					
Dimethylglyoxime	0.258	0.00645	0.0074	0.025	0.258 ± 0.0074
Ni - DTMP -DPG	0.256	0.0107	0,0123	0.042	0.256 ± 0.0123

Table 5. Determination results of nikel (II) in the Sewage water and Bottom sediments ($n = 6$, $P = 0.95$)

Analysis object	Added, μg	Found, μg	Found in the sample, $\mu\text{g} / \text{kg}$	S_r
			$\bar{X} \pm \frac{t_p \cdot S}{\sqrt{n}}$	
Sewage water				
Sample 1	2.0	2.55	0.55 ± 0.05	0.10
Sample 2	5.0	6.64	1.64 ± 0.12	0.07
Bottom sediments				
Sample 1	2.0	3.46	1.46 ± 0.07	0.08
Sample 2	5.0	6.25	2.25 ± 0.04	0.09

4. Conclusions

Nikel(II) forms well chloroform-extractable ternary complex with 2,6-dithiol-4-methylphenol and diphenilguanidine. The

complex could be regarded as an ion associate between an intensively colored anion, $[\text{Ni}(\text{DTMP})_2]^{2-}$, in which nikel is in Ni^{2+} , and a bulky hydrophobic cation (DPGH^+). The

following equilibrium constants and analytical characteristics were determined: constant of extraction, constant of association, constant of distribution, recovery factor, molar absorptivity, Sandell's sensitivity, limit of detection, and limit of quantification. The obtained values show that the studied extraction system in the present work could compete successfully with many similar systems used for nickel determination.

References

- [1] E. Merian, M. Anke, M. Stoppler, Wiley-VCH: Weinheim; Vol.2, (2004). Elements and Their Compounds in the Environment
- [2] C. Ravichandran, D. Benzil, C. Ramachandraiah, Chandrasekhar K. B. Extraction and spectrophotometric determination of nickel in water, alloys and edible oil samples *International Journal of Bioassays* 4.11 (2015): 4468-4472
- [3] H. Avci, G. Kaya, I. Akdeniz, M. Ince and Yaman, M. Flame atomic absorption spectrometric determination of nickel and chromium in various plant leaves used as biomonitors in environmental pollution. *Fresenius Environmental Bulletin*, 2013, 22(2), pp. 379-387.
- [4] R. Dobrowolski, M. Otto. Determination of nickel and cobalt in reference plant materials by carbon slurry sampling GFAAS technique after their simultaneous preconcentration onto modified activated carbon. *Journal of Food Composition and Analysis*, 2012, 26(1-2), pp. 58-65
- [5] O. Sadeghi, N. Tavassoli, M. M. Amini, H. Ebrahimzadeh, N. Daei, Pyridine-functionalized mesoporous silica as an adsorbent material for the determination of nickel and lead in vegetables grown in close proximity by electrothermal atomic adsorption spectroscopy. *Food Chemistry*, 2011, 127(1), pp. 364-368
- [6] C. Zeng, Y. Jia, Y. I. Lee, X. Hou, and L. Wu. Ultrasensitive determination of cobalt and nickel by atomic fluorescence spectrometry using APDC enhanced chemical vapor generation. *Microchemical Journal*. 2012, 104, pp. 33-37
- [7] A. Beiraghi, S. Babae, M. Roshdi. Simultaneous preconcentration of cadmium, cobalt and nickel in water samples by cationic micellar precipitation and their determination by inductively coupled plasma-optical emission spectrometry. *Microchemical Journal*, 2012, 100, pp. 66-71
- [8] K. Uesugi, and S. Yamaguchi. Extraction-Spectrophotometric determination of nickel with 2-hydroxy-1-naphthaldoxime. *Microchemical Journal*, 1982, 27(1), pp. 71-76.
- [9] H. Ishii, T. Odashima, T. Hashimoto. Synthesis of sensitive pyridylhydrazone reagents and extraction-spectrophotometric determination of trace nickel with 2-pyridinecarbaldehyde 2-(5-nitro) pyridylhydrazone. *Analytical Sciences*, 1987. 3(4), pp. 347-352.;
- [10] X. Fan, and C. Zhu, Syntheses of three new benzoic acid-type thiazolylazo reagents and their application to spectrophotometric determination of microamounts of nickel. *Microchemical Journal*, 1998, 59(2), pp. 284-293.;
- [11] N. Chimpalee, D. Chimpalee, P. Keawpasert and D. Thorburn Burns. Flow injection extraction spectrophotometric determination of nickel using bis(acetylacetonate) ethylenediimine. *Analytica Chimica Acta*, 2000, 408(1-2), pp. 123-127.;
- [12] A. K. Malik and A. L. J. Rao. Spectrophotometric determination of cobalt, nickel, palladium, copper, ruthenium and molybdenum after extraction of their isoamyl xanthate complexes into molten naphthalene. *Journal of Analytical Chemistry*, 2000, V.55, No 8, pp. 746-749.
- [13] T. Suresh, S. S. Kumar, N. M. Kottureshawara et al. Spectrophotometric Study of Nitrogen Base Adducts of Nickel(II)-4-methyl-8-quinolinate. *E-Journal of Chemistry* 2008, V. 5, No.2, pp. 404-408.
- [14] F. Umland, A. Janssen, D. Thierig, G. Wunsch. Theorie und praktische anwendung von complexbildnern. Frankfurt am Main, Akademische Verlagsgesellschaft. 1971.

- [15] Z. Marczenko, and M. Balcerzak. Metod'y spektrofotometrii v UF I vidimoy oblastiakh v neorganicheskom analize. Moskow, Binom. Laboratoriya znaniy. 2007.
- [16] K. A. Kuliyeu, N. A. Verdizadeh. Spectroscopic Investigation of the Complex Formation of Niobium Using 2,6-Dithiolphenol and Aminophenols. American Journal of Analytical Chemistry, 2015, 6, pp.746-756.
- [17] K. A. Kuliyeu, N. A. Verdizadeh, A. B. Gadjieva. Liquid-Liquid Extraction and Spectrophotometric Determination of Molybdenum with 2, 6-Dithiolphenol and its Derivatives in the Presence of Hydrophobic Amines // Chemistry Journal 2015, V. 05, № 3, pp. 45-53
- [18] K. A. Kuliyeu, N. A. Verdizadeh, A. B. Gadjieva, S. A. Mamedova. Spectroscopic investigation complex formation of molybdenum with 2, 6-dithiolphenol and its derivatives in the presence of hydrophobic amines. International Journal of Chemical Studies 2016; 4(3): pp. 42-48
- [19] K. A. Kuliyeu. Spectroscopic Investigation Complex Formation of Vanadium Using 2,6-Dithiol-4-Methylphenol And Hydrophob Amins // Journal of Advances in Chemistry, V. 11, № 4, pp.3487-3497.
- [20] K. A. Kuliyeu, N. A. Verdizadeh. Spectroscopic Investigation Complex Formation of Vanadium Using 2,6-Dithiolphenol and Hydrophob Amins // American Journal of Chemistry, 2015, V. 5, №1, pp. 10-18.
- [21] A. Z. Zalov, N. A. Verdizadeh, R. I. Jamalova. Extraction-photometric determination of niobium (V) with 2-hydroxy-5-bromthiophenol and hydrophob amines. Az. Chim. Journ. № 1. 2011. P. 97-102.
- [22] A. Z. Zalov, G. I. Amanullayeva. Spectrophotometric determination of cobalt (II) in a liquid-liquid extraction system containing 2-hydroxy-5-iodothiophenol and diphenylguanidine. IJRDO-Journal Of Applied Science. V.2 | Issue-7 | July,2016 | Paper-2 17-25
- [23] A. Z. Zalov, K. B. Gavazov. Extractive Spectrophotometric Determination of Nickel with 2-hydroxy-5-iodothiophenol and Diphenylguanidine Chemistry Journal (2014), Vol. 04, Issue 5, 20-25
- [24] A. M. Kuliev, S. R. Aliev, F. N. Mamedov and M. Movsumzade. Synthesis of aminomethyl derivatives of 2-hydroxy-5-tert-alkylthiophenols and their cleavage by thiols. Zhurnal Organicheskoi Khimii, 1976, V. 12, No 2, pp. 426-431.
- [25] M. I. Bulatov, I. P. Kalinkin. Prakticheskoe rukovodstvo po photocolorimetriceskim i spectrophotometriceskim metodam analiza, Leningrad, 1976.
- [26] K. Nakamoto. IR-spectra i KR neorganicheskikh i koordinatsionnykh soedinenii (Infrared and Raman Spectra of Inorganic and Coordination Compounds). Moscow: Mir, 1991. P. 536.
- [27] L. Bellami. Infkrasnie spectri slojnykh molekul (The infra-red spectra of complex molecules). Moscow: Mir, 1991. P. 592.
- [28] M. K. Akhmedly, A. E. Kly'gin, L. I. Ivanova, E. A. Bashirov. On the chemistry of interaction of gallium ions with a number of sulphophthalates. Zhurnal Neorganicheskoi Khimii, 1974. V.19.№8. pp. 2007-2012.
- [29] V. A. Nazarenko. Tr. Komis. po analit. khimii Akad. Nauk SSSR (Proc. Commission on Analytical Chemistry of the USSR's Academy of Sciences), 1969, Moscow: Nauka, vol. 17. p. 22.
- [30] V. A. Nazarenko, E. A. Biryuk. A study of the chemistry of reactions of multivalent element ions with organic reagents. Zh. Anal. Khim., 1967. V.22 № 13, pp.57-64.
- [31] Z. Zhiming, M. Dongsen, Y. Cunxiao. "Mobile equilibrium method for determining composition and stability constant of coordination compounds of the form $MmRn$," *Journal of Rare Earths*, 1997. V. 15, No. 3, pp. 218-219.
- [32] A. Holme, F. J. Langmyhr, "A modified and a new straightline method for determining the composition of weak

- complexes of the form AmB_n ," *Analytica Chimica Acta*, vol. 36, pp. 383–391, 1966.
- [33] E. Harvey, D. L. Manning. "Spectrophotometric methods of establishing empirical formulas of colored complexes in solution," *Journal of the American Chemical Society*, 1950. V. 72, No.10, pp. 4488–4493,
- [34] Ye. N. Dorokhova, G. V. Prokhorova. *Analiticheskaya khimiya (fiziko-khimicheskiye metody analiza)*. M.: Vysshaya shkola, 1991.
- [35] N. Yoshikuni, T. Baba, N. Tsunoda, K. Oguma. Aqueous two-phase extraction of nickel dimethylglyoximate complex and its application to spectrophotometric determination of nickel in stainless steel. *Talanta*. 2005, V. 66, No 1, pp. 40–44.
- [36] C. Ramachandraiah, J. R. Kumar, K. J. Reddy et al. Development of a highly sensitive extractive spectrophotometric method for the determination of nickel(II) from environmental matrices using N-ethyl-3-carbazolecarboxaldehyde-3-thiosemicarbazone. *J Environ Manage* 2008 V. 88, No4, pp.729-736.
- [37] V. A. Jadhav, M. U. Kulkarni 7-Methyl-2-chloroquinoline-3-carbaldehyde thiosemicarbazone as analytical reagent for copper, cobalt and nickel (II). *J. Indian Chem Soc* 1992; 69: pp.287–288.
- [38] M. Otomo, T. Watanabe and M. Moriya. Solvent Extraction and Spectrophotometric Determination of Nickel (II) with Thiazole-2-carbaldehyde 2-Quinolylhydrazone *Analytical Sciences*. 1986, V. 2, No. 6, pp. 549-552.
- [39] L. S. Sarma, J. R. Kumar, K. J. Reddy et al. Development of highly sensitive extractive spectrophotometric determination of nickel (II) in medicinal leaves, soil, industrial effluents and standard alloy samples using pyridoxal-4-phenyl-3-thiosemicarbazone. *Journal of Trace Elements in Medicine and Biology* 22 (2008) pp.285–295
- [40] D. Rekha, J. D. Kumar, B. Jayaraj et al. Nickel (II) Determination by Spectrophotometry Coupled with Preconcentration Technique in Water and Alloy Samples *Bull. Korean Chem. Soc.* 2007, Vol. 28, No. 3 pp. 373-378.